

Virial coefficients and equations of state for mixtures of hard discs, hard spheres and hard hyperspheres

A. Santos*

*Department of Physics, University of Florida,
Gainesville, FL 32611, U.S.A.*

S. B. Yuste†

*Institute for Nonlinear Science and Department of Chemistry, University of California,
San Diego, La Jolla, CA 92093-0340, U.S.A.*

M. López de Haro‡

*Centro de Investigación en Energía, UNAM,
Temixco, Morelos 62580, México
(February 1, 2008)*

The composition-independent virial coefficients of a d -dimensional binary mixture of (additive) hard hyperspheres following from a recent proposal for the equation of state of the mixture [SANTOS, A., YUSTE, S. B., AND LÓPEZ DE HARO, M., 1999, *Molec. Phys.*, **96**, 1] are examined. Good agreement between theoretical estimates and available exact or numerical results is found for $d = 2, 3, 4$ and 5 , except for mixtures whose components are very disparate in size. A slight modification that remedies this deficiency is introduced and the resummation of the associated virial series is carried out, leading to a new proposal for the equation of state. The case of binary hard-sphere mixtures ($d = 3$) is analyzed in some detail.

I. INTRODUCTION

Depending on the nature of the independent variables, it is well known that the full thermodynamic description of a given system requires the availability of either one of the thermodynamic potentials or of the corresponding equations of state. In the case of classical fluids, the usual expression for the equation of state (EOS) is a relationship among the pressure p , the density ρ and the temperature T of the fluid. However, except for very few ideal systems, the explicit (exact) form of this relationship is not known in general. Therefore, given the fact that an experimental determination of the EOS for every particular fluid is of course not practical, researchers have concentrated their efforts in proposing “reasonable” empirical or semi-empirical approximations. Perhaps the neatest example of such an approach, already produced over a century ago and providing an essentially correct qualitative picture of the thermodynamic properties of gases and liquids, is the celebrated van der Waals EOS [1]. An alternative proposal, first introduced by Kammerlingh Onnes [2] as a mathematical representation of experimental results on the equation of state of gases and liquids, is the so called virial equation of state. Written for the compressibility factor $Z \equiv p/\rho k_B T$ (where k_B is the Boltzmann constant) as a power series in ρ , it provides an expansion that gives the deviation from ideal gas behaviour in ascending powers of the density, namely

$$Z = 1 + \sum_{n=2}^{\infty} B_n \rho^{n-1}. \quad (1)$$

Here, the B_n are the virial coefficients. For a simple fluid, the virial coefficients are in general functions of T alone, while in the case of mixtures they also depend on composition [3]. An interesting aspect of the virial equation of state, discovered many years after it was introduced, is the fact that it can be derived rigorously using statistical mechanics [4]. This in turn implies that the virial coefficients are not merely empirical constants but are rather related to intermolecular interactions in a well-defined manner. It is unfortunate, however, that in general the actual computation of the virial coefficients is a formidable task and that the radius of convergence of the series in Eq. (1) is not known.

In the case of hard-core fluids (rods, discs, spheres and hyperspheres) the virial coefficients are easier to compute than those corresponding to any other intermolecular interactions. Even in these model systems and with the exception of hard rods, where the exact EOS and corresponding virial coefficients are known, only the first few of such coefficients are available so far [5–8] because the number of cluster integrals involved increases very rapidly with the order of the coefficient. The availability is scarcer for mixtures than for simple fluids, although recently there has been renewed interest in partially remedying this deficiency [9–19]. The limited knowledge of the virial coefficients has proved useful,

however, in the sense that proposals for the EOS of these systems may be judged amongst other things by how well the virial coefficients arising in such proposals compare with the exact values. The rationale here is that a “good” theoretical EOS should lead to an accurate prediction of the value of the virial coefficients. Alternatively, the few available virial coefficients may also be profitably used to construct a rational (Padé or Levin) [6,20–24] approximation to the EOS or to guide in the construction of a theoretical EOS, as was the case with the Carnahan-Starling (CS) EOS [25] for a simple hard-sphere fluid.

A variety of (approximate) EOS for simple fluids composed of hard discs and hard spheres are available in the literature [26] and some work has also been reported on simple fluids of hard hyperspheres [27–31]. Very recently we introduced a simple recipe to derive the compressibility factor of a multicomponent mixture of d -dimensional hard hyperspheres in terms of that of the corresponding single component system [32]. A straightforward consequence of such a recipe is that one can easily derive the explicit expressions for the virial coefficients of the mixture in terms of those of the single component fluid for all d . The major aim of this paper is to further assess the usefulness of our recipe by comparing the predictions of the values of the virial coefficients of binary mixtures of hard discs, hard spheres and hard hyperspheres in $d = 4$ and $d = 5$ with those obtained through Monte Carlo integration. Along the way we will introduce slight modifications to our original proposal and consider an alternative EOS to cope with mixtures very disparate in size. This new EOS is consistent with the forms suggested by Wheatley for the virial coefficients of hard discs [33] and hard spheres [34].

The paper is organized as follows. In the next Section and in order to make the paper self-contained, we will briefly recall the main results of our previous work and write down explicitly the composition-independent virial coefficients of a d -dimensional binary mixture of (additive) hard hyperspheres. Section III deals with a new proposal for the EOS of the binary mixture that attempts to incorporate additional features not present in the original recipe. In Section IV we focus on the case of hard spheres because of its intrinsic interest. We close the paper in Section V with an assessment of the different results. Some mathematical details of our derivations are presented in two Appendices.

II. APPROXIMATE EQUATION OF STATE AND VIRIAL COEFFICIENTS FOR A BINARY MIXTURE OF d -DIMENSIONAL HARD SPHERES

In this Section we first consider a multicomponent mixture of d -dimensional (additive) hard hyperspheres and then restrict ourselves to *binary* mixtures of hard hyperspheres in d dimensions.

Let the number of components be N , the total number density of the mixture be ρ , the set of mole fractions be $\{x_i\}$, and the set of diameters be $\{\sigma_i\}$, ($i = 1, 2, \dots, N$). The packing fraction is $\eta = \sum_{i=1}^N \eta_i = v_d \rho \langle \sigma^d \rangle$, where $\eta_i = v_d \rho_i \sigma_i^d$ is the partial packing fraction due to species i , $\rho_i = \rho x_i$ is the partial number density corresponding to species i , $v_d = (\pi/4)^{d/2} / \Gamma(1 + d/2)$ is the volume of a d -dimensional sphere of unit diameter and $\langle \sigma^n \rangle \equiv \sum_{i=1}^N x_i \sigma_i^n$. In previous work [32] we proposed a simple equation of state for the mixture, $Z_m(\eta)$, consistent with a given EOS for a single component system, $Z_s(\eta)$, at the same packing fraction η , namely

$$Z_m(\eta) = 1 + [Z_s(\eta) - 1] 2^{1-d} \Delta_0 + \frac{\eta}{1 - \eta} \left(1 - \Delta_0 + \frac{1}{2} \Delta_1 \right), \quad (2)$$

with

$$\Delta_p \equiv \frac{\langle \sigma^{d+p-1} \rangle}{\langle \sigma^d \rangle^2} \sum_{m=p}^{d-1} \frac{(d+p-1)!}{m!(d+p-1-m)!} \langle \sigma^{m-p+1} \rangle \langle \sigma^{d-m} \rangle. \quad (3)$$

In the one-dimensional case, Eq. (2) yields the *exact* result $Z_m(\eta) = Z_s(\eta)$. Further, for binary mixtures in $d = 2, 3, 4$ and 5 , it proved to be very satisfactory when a reasonably accurate $Z_s(\eta)$ was taken [24,32,35,36].

From the virial expansion $Z_s(\eta) = 1 + \sum_{n=2}^{\infty} b_n \eta^{n-1}$, where b_n are (reduced) virial coefficients, and according to Eqs. (1)–(3), the n -th virial coefficient of the mixture is given by

$$B_n = v_d^{n-1} \langle \sigma^d \rangle^{n-1} \left(2^{1-d} \Delta_0 b_n + 1 - \Delta_0 + \frac{1}{2} \Delta_1 \right), \quad (4)$$

and explicitly for $n \geq 2$

$$B_n = v_d^{n-1} \langle \sigma^d \rangle^{n-3} \sum_{m=0}^d \frac{(d-1)!}{m!(d-m)!} \langle \sigma^{d-m} \rangle \left[(2^{1-d} b_n - 1)(d-m) \langle \sigma^{m+1} \rangle \langle \sigma^{d-1} \rangle + \frac{d}{2} \langle \sigma^m \rangle \langle \sigma^d \rangle \right]. \quad (5)$$

Now we particularize to *binary* mixtures. In that case, it is useful to define composition-independent coefficients B_{n_1, n_2} as

$$B_n = \sum_{n_1=0}^n B_{n_1, n-n_1} \frac{n!}{n_1!(n-n_1)!} x_1^{n_1} x_2^{n-n_1}. \quad (6)$$

Our objective is to use our model to get approximate expressions of B_{n_1, n_2} as explicit functions of σ_1 , σ_2 , and d . In Appendix A it is shown that the result is

$$B_{n_1, n_2} = v_d^{n-1} \sigma_1^{d(n-1)} \alpha^{d(n_2-1)} \left[C_{n_1, n_2}^{(1)} \alpha^d + C_{n_1, n_2}^{(2)} \alpha^{d-1} + C_{n_1, n_2}^{(3)} (1+\alpha)^{d-1} + C_{n_2, n_1}^{(3)} \alpha (1+\alpha)^{d-1} + C_{n_2, n_1}^{(2)} \alpha + C_{n_2, n_1}^{(1)} \right], \quad (7)$$

where $n \equiv n_1 + n_2$, $\alpha \equiv \sigma_2/\sigma_1$ and

$$C_{n_1, n_2}^{(1)} = \frac{n_1(n_1-1) [(n_1-2)b_n + 2^{d-1}n_2]}{n(n-1)(n-2)}, \quad (8)$$

$$C_{n_1, n_2}^{(2)} = \frac{n_1 n_2 (n_1-1) (b_n - 2^{d-1})}{n(n-1)(n-2)}, \quad (9)$$

$$C_{n_1, n_2}^{(3)} = \frac{n_1 n_2 [(n_2-1)2^{2-d}b_n + n_1 - n_2]}{n(n-1)(n-2)}. \quad (10)$$

The special case of $n = 2$ is obtained by *first* setting $b_n = 2^{d-1}$ and *then* setting $n = 2$. The result is

$$C_{n_1, n_2}^{(1)} = 2^{d-2} n_1 (n_1 - 1), \quad C_{n_1, n_2}^{(2)} = 0, \quad C_{n_1, n_2}^{(3)} = \frac{1}{2} n_1 n_2. \quad (11)$$

This gives the *exact* second virial coefficient.

Equation (7) expresses the reduced coefficient

$$B_{n_1, n_2}^*(\alpha) \equiv B_{n_1, n_2}(\sigma_1, \sigma_2) \sigma_1^{-d(n-1)} \alpha^{-d(n_2-1)} \quad (12)$$

as a polynomial in α of degree d . Although it is known that the *exact* coefficients do not have in general such a polynomial structure [34], most of the proposals to date for $d = 2$ and $d = 3$ are polynomials [15,16,18,23,24,33,34]. Let us see which consistency conditions the approximation (7)–(10) fulfills. First, it verifies the obvious symmetry property $B_{n_1, n_2}(\sigma_1, \sigma_2) = B_{n_2, n_1}(\sigma_2, \sigma_1)$, which implies that $B_{n_1, n_2}^*(\alpha) = \alpha^d B_{n_2, n_1}^*(1/\alpha)$. In addition, the coefficients (8)–(10) satisfy the properties

$$C_{n_1, n_2}^{(1)} + C_{n_2, n_1}^{(2)} + 2^{d-1} C_{n_2, n_1}^{(3)} = \frac{n_1}{n} b_n, \quad (13)$$

$$d C_{n_1, n_2}^{(1)} + (d-2) C_{n_1, n_2}^{(2)} + 2^{d-1} C_{n_2, n_1}^{(3)} = \frac{n_1}{n(n-1)} [2^{d-1} n_2 + d(n_1-1)b_n], \quad (14)$$

$$C_{n, 0}^{(1)} = b_n, \quad C_{0, n}^{(1)} = C_{n, 0}^{(2)} = C_{0, n}^{(2)} = C_{n, 0}^{(3)} = C_{0, n}^{(3)} = 0. \quad (15)$$

Equation (13) implies that

$$C_{n_1, n_2}^{(1)} + C_{n_2, n_1}^{(1)} + C_{n_1, n_2}^{(2)} + C_{n_2, n_1}^{(2)} + 2^{d-1} [C_{n_1, n_2}^{(3)} + C_{n_2, n_1}^{(3)}] = b_n, \quad (16)$$

which guarantees that

$$B_{n_1, n_2}^*(\alpha = 1) = v_d^{n-1} b_n, \quad (17)$$

i.e., if both species have the same size we recover the one-component case. The same situation appears if $x_1 = 0$ or $x_2 = 0$, which means that $B_{n,0}^*(\alpha) = v_d^{n-1} b_n \alpha^d$ and $B_{0,n}^*(\alpha) = v_d^{n-1} b_n$. This is verified as a consequence of Eq. (15). A subtler consistency condition is [34]

$$\left. \frac{\partial B_{n_1, n_2}^*}{\partial \alpha} \right|_{\alpha=1} = d \frac{n_1}{n} v_d^{n-1} b_n, \quad (18)$$

which ensures that the derivative of the excess free energy of mixing with respect to σ_2 is zero when the spheres are of the same size [34]. In the approximation (7), Eq. (18) requires

$$dC_{n_1, n_2}^{(1)} + (d-1)C_{n_1, n_2}^{(2)} + 2^{d-2}(d-1)C_{n_1, n_2}^{(3)} + 2^{d-2}(d+1)C_{n_2, n_1}^{(3)} + C_{n_2, n_1}^{(2)} = d \frac{n_1}{n} b_n. \quad (19)$$

If we interchange the roles of n_1 and n_2 in Eq. (19) and add the original and the transformed equations, we get Eq. (16). If, instead, we subtract both equations, we obtain

$$d \left[C_{n_1, n_2}^{(1)} - C_{n_2, n_1}^{(1)} \right] + (d-2) \left[C_{n_1, n_2}^{(2)} - C_{n_2, n_1}^{(2)} \right] - 2^{d-1} \left[C_{n_1, n_2}^{(3)} - C_{n_2, n_1}^{(3)} \right] = d \frac{n_1 - n_2}{n} b_n. \quad (20)$$

Thus, enforcement of Eq. (19) is equivalent to enforcement of Eq. (20). The property (14) guarantees that the condition (20) is fulfilled by the coefficients (8)–(10).

The most stringent conditions appear in the limits $\alpha \rightarrow 0$ and $\alpha \rightarrow \infty$. In that case, Eq. (7) yields

$$B_{n_1, n_2}^*(\alpha) = \begin{cases} v_d^{n-1} \left[C_{n_1, n_2}^{(3)} + C_{n_2, n_1}^{(1)} \right] & (\alpha \rightarrow 0), \\ v_d^{n-1} \alpha^d \left[C_{n_1, n_2}^{(1)} + C_{n_2, n_1}^{(3)} \right] & (\alpha \rightarrow \infty). \end{cases} \quad (21)$$

On the other hand, the *exact* result is [33,34,37]

$$B_{n_1, n_2}^*(\alpha) = \begin{cases} v_d^{n-1} \frac{n_2}{n} b_{n_2} & (\alpha \rightarrow 0), \\ v_d^{n-1} \alpha^d \frac{n_1}{n} b_{n_1} & (\alpha \rightarrow \infty). \end{cases} \quad (22)$$

The above condition means that in the limit of infinite size asymmetry, the smaller spheres contribute to the total pressure as if they were a one-component system in a free volume equal to the total volume minus the volume occupied by the larger spheres. Our approximation (7)–(10) gives the correct forms for the asymptotic behaviours of B_{n_1, n_2}^* , but not the correct coefficients (except for $n_2 = 1$ in the limit $\alpha \rightarrow 0$ and for $n_1 = 1$ in the limit $\alpha \rightarrow \infty$, apart from the trivial cases $n_1 = 0, n$ and $n_2 = 0, n$). In fact, in our approximation the virial coefficient B_n depends parametrically on b_n only, and not on the previous coefficients $b_{n_1}, n_1 \leq n$, as is needed in Eq. (22). This is a consequence of the form of our approximation in which the compressibility factor of the mixture at a given packing fraction η is expressed in terms of the compressibility factor of the monodisperse system at the *same* packing fraction.

Another consistency condition is [34]

$$\left. \frac{\partial B_{n-1, 1}^*}{\partial \alpha} \right|_{\alpha=0} = v_d^{n-1} d \frac{n-1}{n}. \quad (23)$$

This comes from the exact value of the derivative of the excess free energy with respect to molecular size in the limit of infinite size asymmetry [38]. In general, this condition is not verified by our approximation either. We consider the condition (22) to be more important than (23) because it is not restricted to $n_2 = 1$ (or $n_1 = 1$) and also because it refers to the leading term in the limit of infinite size asymmetry, while the condition (23) refers to the sub-leading term.

TABLE I. Reduced virial coefficients b_n for the one-component case.

n	$d = 2^a$	$d = 3^a$	$d = 4^b$	$d = 5^b$
2	2	4	8	16
3	3.12801775	10	32.4057594	106
4	4.25785446	18.36477	77.7451797	311.18341
5	5.336897	28.2245	145.9	843.4
6	6.3626	39.739	252.0	988
7	7.351	53.539		
8	8.338	70.78		

^aRef. [6]

^bRefs. [7], [29] and [30]

Table I gives the known values of the coefficients b_n for $2 \leq d \leq 5$. From them, Eqs. (7)–(11) can be used to estimate the coefficients B_{n_1, n_2}^* for any value of α in these dimensions. We will come back to this point later on. The values of Table I can also be used to check to which degree Eq. (22) is violated by the approximation (7)–(10). The less favorable case corresponds to $n_1 = 1$ in the limit $\alpha \rightarrow 0$ (or, equivalently, $n_2 = 1$ in the limit $\alpha \rightarrow \infty$). Let us then define

$$R_n \equiv \frac{C_{1, n-1}^{(3)} + C_{n-1, 1}^{(1)}}{[(n-1)/n]b_{n-1}}. \quad (24)$$

According to Eqs. (8) and (10), we have

$$R_n = \frac{(2^{2-d} + n - 3)b_n + 2^{d-1} - 1}{(n-1)b_{n-1}}. \quad (25)$$

Table II shows the values of R_n that can be obtained from the known virial coefficients. In the cases $d = 2$ and $d = 3$ the known values of the ratio R_n differ from 1 less than 5%. In fact, by assuming $R_n \simeq 1$, Eq. (25) can be used to estimate b_9 and b_{10} . This gives $b_9 \simeq 9.39$ ($d = 2$), $b_9 \simeq 86.7$ ($d = 3$), $b_{10} \simeq 10.44$ ($d = 2$) and $b_{10} \simeq 103.6$ ($d = 3$). Estimates based on Padé approximants are [6] $b_9 \simeq 9.37$ ($d = 2$), $b_9 \simeq 93.1$ ($d = 3$), $b_{10} \simeq 10.55$ ($d = 2$) and $b_{10} \simeq 123.2$ ($d = 3$). On the other hand, the deviations of R_n from 1 become more important as the dimensionality increases and are already relatively large for $d = 5$. Thus, at least for those high dimensionalities, it appears to be convenient to modify the prescription (7) as to satisfy the requirement (22). This is the subject of the next Section, where we will also discuss in more detail how good the estimates of B_{n_1, n_2}^* , as obtained after using Eqs. (7)–(11) with the values of the coefficients b_n given in Table I, are for any value of α and $2 \leq d \leq 5$.

TABLE II. Ratio R_n , as given by Eq. (25).

n	$d = 2$	$d = 3$	$d = 4$	$d = 5$
3	1.032	1	0.944	0.883
4	1.014	1.018	1.072	1.148
5	0.999	1.001	1.078	1.452
6	0.991	1.007	1.132	0.736
7	0.989	1.023		
8	0.992	1.047		

III. A NEW PROPOSAL FOR THE EQUATION OF STATE OF A BINARY MIXTURE

In order to account for the conditions imposed by Eq. (22) without greatly sacrificing the simplicity of our original recipe, let us now explore the possibility of keeping the structure (7), but with expressions for the coefficients different from (8)–(10). The requirement (22) implies that

$$C_{n_1, n_2}^{(3)} + C_{n_2, n_1}^{(1)} = \frac{n_2}{n} b_{n_2}. \quad (26)$$

In addition, the condition (17) yields Eq. (16). Note that Eq. (13) is more restrictive than Eq. (16). Since we want our modified expressions to remain as close as possible to the original ones, we will take (13) as a second condition. From Eqs. (26) and (13) we have

$$C_{n_1, n_2}^{(1)} = (2^{d-1} - 1)^{-1} \left[\frac{n_1}{n} (2^{d-1} b_{n_1} - b_n) + C_{n_2, n_1}^{(2)} \right], \quad (27)$$

$$C_{n_1, n_2}^{(3)} = (2^{d-1} - 1)^{-1} \left[\frac{n_2}{n} (b_n - b_{n_2}) - C_{n_1, n_2}^{(2)} \right]. \quad (28)$$

We need a third condition to close the problem. Since $C_{n_1, n_2}^{(2)}$ does not enter into Eq. (21) and in agreement with our philosophy of departing from the original formulation as less as possible, one possibility is to keep Eq. (9); however, the resulting approximation is consistent with the conditions (15) but does not satisfy the requirement (19). If we enforce the fulfillment of Eq. (19) [or, equivalently, of Eq. (20)], then $C_{n_1, n_2}^{(2)}$ must have the following form

$$C_{n_1, n_2}^{(2)} = D_{n_1, n_2} - \frac{n_1}{n} \frac{b_n - b_{n_1}}{2^{2-d} - 1}, \quad (29)$$

where D_{n_1, n_2} must be *symmetric*, but otherwise remains so far unknown. To close the problem, we impose the more restrictive property (14). This gives rise to

$$D_{n_1, n_2} = \frac{1}{n [2 + 2^{d-1}(d-3)]} \left[(2^{d-1} - 1) (2^{d-1} - d b_n) \frac{n_1 n_2}{n - 1} + \frac{2^{d-1} - d}{1 - 2^{2-d}} (n b_n - n_1 b_{n_1} - n_2 b_{n_2}) \right]. \quad (30)$$

Equations (27)–(30) close the modified version of our approximation.

It must be noted that Eqs. (29) and (30) are meaningless if $d = 2$. This is related to the fact that if $d = 2$ it is impossible to enforce conditions (17), (18) and (22) simultaneously on a polynomial of second degree for B_{n_1, n_2}^* , as the number of conditions exceeds the number of unknowns. As a matter of fact, Eqs. (27) and (28) are enough to uniquely determine B_{n_1, n_2}^* in the case $d = 2$, irrespective of the value of $C_{n_1, n_2}^{(2)}$. The result is

$$B_{n_1, n_2}^* = v_2^{n-1} \left[\frac{n_2}{n} b_{n_2} + \left(b_n - \frac{n_1}{n} b_{n_1} - \frac{n_2}{n} b_{n_2} \right) \alpha + \frac{n_1}{n} b_{n_1} \alpha^2 \right]. \quad (31)$$

This was the form proposed by Wheatley [33] by imposing conditions (17) and (22). However, condition (18) is not verified unless $n_1(b_n - b_{n_1}) = n_2(b_n - b_{n_2})$, which is only true if $b_n = n$, i.e., if one considers the scaled particle theory [39]. For the sake of comparison, the form that stems from our original approximation, Eqs. (7)–(10), is (for $d = 2$)

$$B_{n_1, n_2}^* = v_2^{n-1} \frac{n_1 n_2}{n(n-1)} \left[\frac{n_2 - 1}{n_1} b_n + 1 + 2(b_n - 1)\alpha + \left(\frac{n_1 - 1}{n_2} b_n + 1 \right) \alpha^2 \right]. \quad (32)$$

In summary, the modified expressions for the virial coefficients are given by Eq. (31) for $d = 2$ and by Eqs. (7) and (27)–(30) for $d \geq 3$. We are now in a position to compare the theoretical predictions we have provided in this and the previous Section with the available Monte Carlo data for the virial coefficients. We note that the coefficients B_{n_1, n_2} corresponding to $n \leq 5$ have been recently evaluated for $d = 2$ [11,15], $d = 3$ [12–14,16], $d = 4$ [19] and $d = 5$ [19]. Further, the coefficients corresponding to $n = 6$ for $d = 2$ are also known [18]. Because of its special physical interest and for reasons that will become apparent later, the case of hard spheres ($d = 3$) will be addressed separately in Section IV.

We begin our assessment of our estimates of B_{n_1, n_2} with the binary mixture of hard discs. This implies using Eq. (32) for the original recipe or using Eq. (31) in the modified version, which coincides with Wheatley's proposal [33]. In Figs. 1–4 we compare the results of taking these two routes with the ones of Refs. [10,11,15,18]. As clearly seen in the figures, the theoretical prescriptions do a very good job for the whole range of α , the modified version (31) being almost perfect.




FIG. 1. Reduced composition-independent virial coefficients $B_{n_1, n-n_1}^*$ as functions of the size ratio α for $d = 2$, $n = 3$, and, from top to bottom, $n_1 = 1$ and 2. The circles are exact results [10], the solid line is the theoretical prediction (32) and the dashed line is the theoretical prediction (31).




FIG. 2. Reduced composition-independent virial coefficients $B_{n_1, n-n_1}^*$ as functions of the size ratio α for $d = 2$, $n = 4$, and, from top to bottom, $n_1 = 1, 2$ and 3. The circles are “exact” Monte Carlo results [11], the solid line is the theoretical prediction (32) and the dashed line is the theoretical prediction (31).




FIG. 3. Reduced composition-independent virial coefficients $B_{n_1, n-n_1}^*$ as functions of the size ratio α for $d = 2$, $n = 5$, and, from top to bottom, $n_1 = 1, 2, 3$ and 4 . The circles are “exact” Monte Carlo results [15], the solid line is the theoretical prediction (32) and the dashed line is the theoretical prediction (31).




FIG. 4. Reduced composition-independent virial coefficients $B_{n_1, n-n_1}^*$ as functions of the size ratio α for $d = 2$, $n = 6$, and, from top to bottom, $n_1 = 1, 2, 3, 4$ and 5 . The circles are “exact” Monte Carlo results [18], the solid line is the theoretical prediction (32) and the dashed line is the theoretical prediction (31).

In the case of hard hyperspheres in $d = 4$ and $d = 5$, Figs. 5–10 illustrate the performance of Eq. (7) together with (8)–(11) and of Eqs. (7) and (27)–(30) with respect to the “exact” data of Ref. [19]. In these dimensionalities, it is clear that while the modified version of the estimates for the virial coefficients continues to exhibit a very good performance, the use of Eqs. (7)–(11) presents limitations for $n = 4$ and especially for $n = 5$. Nevertheless, the EOS for the binary mixture derived from Eq. (2) (which yields the original estimates of the coefficients B_{n_1, n_2}), once the proper $Z_s(\eta)$ is used in both $d = 4$ and $d = 5$, turns out to be very accurate [36].




FIG. 5. Reduced composition-independent virial coefficients $B^*_{n_1, n-n_1}$ as functions of the size ratio α for $d = 4$, $n = 3$, and, from top to bottom, $n_1 = 1$ and 2. The circles are “exact” Monte Carlo results [19], the solid line is the theoretical prediction (7)–(10) and the dashed line is the theoretical prediction given by Eqs. (7) and (27)–(30).




FIG. 6. Reduced composition-independent virial coefficients $B_{n_1, n-n_1}^*$ as functions of the size ratio α for $d = 4$, $n = 4$, and, from top to bottom, $n_1 = 1, 2$ and 3 . The circles are “exact” Monte Carlo results [19], the solid line is the theoretical prediction (7)–(10) and the dashed line is the theoretical prediction given by Eqs. (7) and (27)–(30).




FIG. 7. Reduced composition-independent virial coefficients $B_{n_1, n-n_1}^*$ as functions of the size ratio α for $d = 4$, $n = 5$, and, from top to bottom, $n_1 = 1, 2, 3$ and 4 . The circles are “exact” Monte Carlo results [19], the solid line is the theoretical prediction (7)–(10) and the dashed line is the theoretical prediction given by Eqs. (7) and (27)–(30).




FIG. 8. Reduced composition-independent virial coefficients $B_{n_1, n-n_1}^*$ as functions of the size ratio α for $d = 5$, $n = 3$, and, from top to bottom, $n_1 = 1$ and 2. The circles are “exact” Monte Carlo results [19], the solid line is the theoretical prediction (7)–(10) and the dashed line is the theoretical prediction given by Eqs. (7) and (27)–(30).




FIG. 9. Reduced composition-independent virial coefficients $B_{n_1, n-n_1}^*$ as functions of the size ratio α for $d = 5$, $n = 4$, and, from top to bottom, $n_1 = 1, 2$ and 3. The circles are “exact” Monte Carlo results [19], the solid line is the theoretical prediction (7)–(10) and the dashed line is the theoretical prediction given by Eqs. (7) and (27)–(30).




FIG. 10. Reduced composition-independent virial coefficients $B_{n_1, n-n_1}^*$ as functions of the size ratio α for $d = 5$, $n = 5$, and, from top to bottom, $n_1 = 1, 2, 3$ and 4 . The circles are “exact” Monte Carlo results [19], the solid line is the theoretical prediction (7)–(10) and the dashed line is the theoretical prediction given by Eqs. (7) and (27)–(30).

Once we have analyzed the behaviour of the virial coefficients, the important question that now arises is, which is the EOS consistent with the modified virial coefficients? In the case $d = 2$, from Eq. (31) one gets [37]

$$Z_m(\eta) = \frac{x_1}{1-\eta_2} Z_s \left(\frac{\eta_1}{1-\eta_2} \right) \frac{\sigma_2 - \sigma_1}{\sigma_2} + \frac{x_2}{1-\eta_1} Z_s \left(\frac{\eta_2}{1-\eta_1} \right) \frac{\sigma_1 - \sigma_2}{\sigma_1} + Z_s(\eta) \frac{\langle \sigma^2 \rangle}{\sigma_1 \sigma_2}. \quad (33)$$

The EOS consistent with the coefficients (27)–(30) can also be explicitly obtained for $d \geq 3$ by carrying out the resummation of the corresponding virial series. The result is (see Appendix B for details)

$$Z_m(\eta) = 1 + \frac{K}{\langle \sigma^d \rangle} [Z_s(\eta) - 1] + \frac{K_0}{\langle \sigma^d \rangle} x_1 x_2 \frac{\eta}{1-\eta} + \frac{K_1}{\sigma_2^d} x_1 \left[\frac{1}{1-\eta_2} Z_s \left(\frac{\eta_1}{1-\eta_2} \right) - 1 \right] + \frac{K_2}{\sigma_1^d} x_2 \left[\frac{1}{1-\eta_1} Z_s \left(\frac{\eta_2}{1-\eta_1} \right) - 1 \right], \quad (34)$$

where K , K_0 , K_1 and K_2 are independent of the density and are given by

$$K = x_1^2 \sigma_1^d \left(1 - \frac{K_1}{\sigma_2^d} \right) + x_2^2 \sigma_2^d \left(1 - \frac{K_2}{\sigma_1^d} \right) + 2x_1 x_2 \left(\sigma_{12}^d - \frac{K_1 + K_2 + K_0}{2^d} \right), \quad (35)$$

$$K_0 = \frac{1}{2^{2-d} + d - 3} [\sigma_1^d + \sigma_2^d + (2^{d-1} - 1) \sigma_1 \sigma_2 (\sigma_1^{d-2} + \sigma_2^{d-2}) - 2^d \sigma_{12}^d], \quad (36)$$

$$K_1 = \frac{1}{(2^d - 4)(2^{2-d} + d - 3)\sigma_{12}} \{ \sigma_{12}^d [2^d(d-2)\sigma_1 + 8(1 - 2^{d-1} + d2^{d-3})\sigma_2] - (2^{d-1} - 2)\sigma_1 \sigma_2 (\sigma_1^{d-1} + \sigma_2^{d-1}) - (d-2)(2^{d-1} - 1)\sigma_1^2 \sigma_2^{d-1} - (2^{d-1} - d)\sigma_1^{d-1} \sigma_2^2 - (d-2)\sigma_1^{d+1} - [d-4-2^{d-1}(d-3)]\sigma_2^{d+1} \}, \quad (37)$$

$$K_2 = \frac{1}{(2^d - 4)(2^{2-d} + d - 3)\sigma_{12}} \left\{ \sigma_{12}^d [2^d(d-2)\sigma_2 - 8(1 - 2^{d-1} + d2^{d-3})\sigma_1] \right. \\ \left. - (2^{d-1} - 2)\sigma_1\sigma_2(\sigma_1^{d-1} + \sigma_2^{d-1}) - (d-2)(2^{d-1} - 1)\sigma_2^2\sigma_1^{d-1} - (2^{d-1} - d)\sigma_2^{d-1}\sigma_1^2 \right. \\ \left. - (d-2)\sigma_2^{d+1} - [d-4-2^{d-1}(d-3)]\sigma_1^{d+1} \right\}. \quad (38)$$

Here, $\sigma_{12} \equiv (\sigma_1 + \sigma_2)/2$. Note that Eq. (33) for hard discs is included in the structure of Eq. (34), except that then Eqs. (35)–(38) cannot be applied and we have instead $K_0 = 0$, $K_1 = \sigma_2(\sigma_2 - \sigma_1)$, $K_2 = \sigma_1(\sigma_1 - \sigma_2)$ and $K = \langle \sigma^2 \rangle (1 - x_1 K_1/\sigma_2^2 - x_2 K_2/\sigma_1^2)$.

Equation (34) complies with the exact results given in Eq. (22). It is clearly more complicated than the original recipe, Eq. (2), and should in principle be useful in particular for mixtures involving components of disparate sizes. In contrast to Eq. (2), the proposal (34) expresses $Z_m(\eta)$ in terms not only of $Z_s(\eta)$ but also involves $Z_s\left(\frac{\eta_1}{1-\eta_2}\right)$ and $Z_s\left(\frac{\eta_2}{1-\eta_1}\right)$.

We note that if $d = 3$, it follows from Eqs. (35)–(38) that $K_0 = 0$, $K_1 = \sigma_2(\sigma_1 - \sigma_2)^2$, $K_2 = \sigma_1(\sigma_1 - \sigma_2)^2$ and $K = \langle \sigma^3 \rangle (1 - x_1 K_1/\sigma_2^3 - x_2 K_2/\sigma_1^3)$. Consequently, Eq. (34) for hard spheres becomes

$$Z_m(\eta) = \frac{x_1}{1-\eta_2} Z_s\left(\frac{\eta_1}{1-\eta_2}\right) \frac{(\sigma_1 - \sigma_2)^2}{\sigma_2^2} + \frac{x_2}{1-\eta_1} Z_s\left(\frac{\eta_2}{1-\eta_1}\right) \frac{(\sigma_1 - \sigma_2)^2}{\sigma_1^2} \\ + Z_s(\eta) \frac{2\langle \sigma^2 \rangle \sigma_1 \sigma_2 - \langle \sigma^4 \rangle}{\sigma_1^2 \sigma_2^2}. \quad (39)$$

It is interesting to note that Eqs. (33) and (39) can be written in the common form

$$Z_m(\eta) = Z_s(\eta) + x_1 \left[\frac{1}{1-\eta_2} Z_s\left(\frac{\eta_1}{1-\eta_2}\right) - Z_s(\eta) \right] \left(\frac{\sigma_2 - \sigma_1}{\sigma_2} \right)^{d-1} \\ + x_2 \left[\frac{1}{1-\eta_1} Z_s\left(\frac{\eta_2}{1-\eta_1}\right) - Z_s(\eta) \right] \left(\frac{\sigma_1 - \sigma_2}{\sigma_1} \right)^{d-1}. \quad (40)$$

Equation (40) reduces to Eq. (33) if $d = 2$ and to Eq. (39) if $d = 3$. In addition, it becomes exact for $d = 1$, i.e., $Z_s(\eta) = Z_m(\eta) = (1-\eta)^{-1}$. It seems then very tempting to assume Eq. (40) for arbitrary d . With such an assumption, after using Eqs. (B3), (B4), (B8) and (B9), we simply get for the composition-independent virial coefficients

$$B_{n_1, n_2}^* = v_d^{n-1} \left[b_n \frac{n_1 \alpha^d + n_2}{n} + (b_{n_1} - b_n) \frac{n_1}{n} \alpha(\alpha-1)^{d-1} + (b_{n_2} - b_n) \frac{n_2}{n} (1-\alpha)^{d-1} \right]. \quad (41)$$

Of course, this includes the cases $d = 2$ [cf. Eq. (31)] and $d = 3$ [cf. Eq. (47)]. However, Eq. (41) differs from the prescription (27)–(30) if $d \geq 4$. Although Eq. (41) is consistent with conditions (17), (18) [for $d \geq 3$] and (22), it does not reproduce the exact second virial coefficient $B_{1,1}^*(\alpha) = v_d(1+\alpha)^d/2$ (except, of course, for $d \leq 3$). In fact, we have checked that the performance of Eqs. (40) and (41) is very poor for $d \geq 4$, so that they will not be further considered here.

IV. THE EQUATION OF STATE OF BINARY HARD-SPHERE MIXTURES

Up to here, although occasionally we have mentioned explicit features of hard discs and hard spheres, the developments presented in the previous Sections apply for general dimensionality d . Given their intrinsic importance, in this Section we focus on the results for the EOS in three dimensions. In order to set the framework of our discussion, we shall first recall two other EOS in the literature which share with ours the idea of obtaining $Z_m(\eta)$ from the knowledge of $Z_s(\eta)$.

The first one is due to Hamad [40] who in 1994 [41] derived a consistency condition involving the derivatives of the contact values of the radial distribution functions g_{ij} with respect to σ_k [42]. Imposing this condition, he proposed the following equation of state for hard-sphere mixtures

$$Z_m^H(\eta) = Z_s(\eta) + \frac{3\eta}{(1-\eta)^3} \left[\frac{\langle \sigma^2 \rangle^3}{\langle \sigma^3 \rangle^2} \eta + \frac{\langle \sigma \rangle \langle \sigma^2 \rangle}{\langle \sigma^3 \rangle} (1-\eta) - 1 \right], \quad (42)$$

where we have used the label H to refer to Hamad's proposal. It must be noted that Eq. (42) was not explicitly written in Ref. [40], where the results are expressed in a rather more involved form. The second EOS for binary mixtures of hard spheres that we want to consider was proposed more recently by Barrio and Solana [43] (label BS). It reads

$$Z_m^{\text{BS}}(\eta) = 1 + \frac{1}{4}(1 + \beta\eta) \left(1 + 3 \frac{\langle \sigma \rangle \langle \sigma^2 \rangle}{\langle \sigma^3 \rangle} \right) [Z_s(\eta) - 1], \quad (43)$$

where β is adjusted as to reproduce the exact third virial coefficient [9], namely

$$\begin{aligned} B_{3,0}^* &= v_3^2 10 \alpha^3, & B_{2,1}^* &= v_3^2 \left(\frac{1}{3} + 2\alpha + 5\alpha^2 + \frac{8}{3}\alpha^3 \right), \\ B_{1,2}^* &= v_3^2 \left(\frac{8}{3} + 5\alpha + 2\alpha^2 + \frac{1}{3}\alpha^3 \right), & B_{0,3}^* &= v_3^2 10. \end{aligned} \quad (44)$$

More explicitly, in our notation β is written as

$$\beta = \frac{B_3}{v_3^2 \langle \sigma^3 \rangle^2 (1 + 3 \langle \sigma \rangle \langle \sigma^2 \rangle / \langle \sigma^3 \rangle)} - \frac{10}{4}. \quad (45)$$

We now first concentrate on the composition-independent virial coefficients. Our original recipe, Eqs. (7)–(10), gives for $d = 3$

$$\begin{aligned} B_{n_1, n_2}^{\text{SYH}} &= v_3^{n-1} \frac{n_1 n_2}{2n(n-1)(n-2)} \left\{ 2(n_1 + 3n_2 - 4) + \frac{n_2 - 1}{n_1} (n_1 + 2n_2 - 4) b_n \right. \\ &\quad + [2(n_1 - 5n_2 + 4) + (n_1 + 4n_2 - 5) b_n] \alpha + [2(n_2 - 5n_1 + 4) + (n_2 + 4n_1 - 5) b_n] \alpha^2 \\ &\quad \left. + \left[2(n_2 + 3n_1 - 4) + \frac{n_1 - 1}{n_2} (n_2 + 2n_1 - 4) b_n \right] \alpha^3 \right\}, \end{aligned} \quad (46)$$

where we have incorporated the label SYH to distinguish it from the other results. On the other hand, in the three-dimensional case, Eqs. (27)–(30) yield

$$\begin{aligned} B_{n_1, n_2}^{*W} &= v_3^{n-1} \left[\frac{n_2}{n} b_{n_2} + \left(\frac{2n_2 - n_1}{n} b_n + \frac{n_1}{n} b_{n_1} - 2 \frac{n_2}{n} b_{n_2} \right) \alpha \right. \\ &\quad \left. + \left(\frac{2n_1 - n_2}{n} b_n + \frac{n_2}{n} b_{n_2} - 2 \frac{n_1}{n} b_{n_1} \right) \alpha^2 + \frac{n_1}{n} b_{n_1} \alpha^3 \right], \end{aligned} \quad (47)$$

which coincides with the form proposed by Wheatley [34] and so the label in this case is W. The virial coefficients associated with Eqs. (42) and (43) are also readily derived. In the former case, first it is easy to obtain

$$B_n^{\text{H}} = v_3^{n-1} \langle \sigma^3 \rangle^{n-1} \left[b_n - \frac{3}{2} n(n-1) + 3(n-1) \frac{\langle \sigma \rangle \langle \sigma^2 \rangle}{\langle \sigma^3 \rangle} + \frac{3}{2} (n-1)(n-2) \frac{\langle \sigma^2 \rangle^3}{\langle \sigma^3 \rangle^2} \right]. \quad (48)$$

Now, by making $m = 0$ and $m = 1$ in Eq. (A2) and $m = 1$ in Eq. (A1), and inserting the results into Eq. (48), we get the composition-independent coefficients arising from Hamad's proposal,

$$\begin{aligned} B_{n_1, n_2}^{*H} &= v_3^{n-1} \left\{ \frac{n_2}{n} \left[b_n - \frac{3}{2} n_1(n-1+n_2) \right] + \frac{3}{2} \frac{n_1 n_2}{n} (3n_2 - 1) \alpha \right. \\ &\quad \left. + \frac{3}{2} \frac{n_1 n_2}{n} (3n_1 - 1) \alpha^2 + \frac{n_1}{n} \left[b_n - \frac{3}{2} n_2(n-1+n_1) \right] \alpha^3 \right\}. \end{aligned} \quad (49)$$

As far as Eq. (43) is concerned, one readily derives (for $n \geq 3$)

$$\begin{aligned} B_n^{\text{BS}} &= \frac{v_3^{n-1}}{4} (b_n + \beta b_{n-1}) \langle \sigma^3 \rangle^{n-1} \left(1 + 3 \frac{\langle \sigma \rangle \langle \sigma^2 \rangle}{\langle \sigma^3 \rangle} \right) \\ &= \frac{v_3^{n-1}}{4} \left(b_n - \frac{10}{4} b_{n-1} \right) \langle \sigma^3 \rangle^{n-1} \left(1 + 3 \frac{\langle \sigma \rangle \langle \sigma^2 \rangle}{\langle \sigma^3 \rangle} \right) + \frac{v_3^{n-3}}{4} b_{n-1} B_3 \langle \sigma^3 \rangle^{n-3}. \end{aligned} \quad (50)$$

Now, after some algebra also involving the substitution of the results obtained by taking $m = 0$ and $m = 1$ in Eq. (A2) into Eq. (50), the composition-independent virial coefficients for the Barrio-Solana EOS turn out to be given by

$$\begin{aligned}
B_{n_1, n_2}^{*BS} = & v_3^{n-1} \frac{b_n}{4n(n-1)} [n_2(3n_2 + n - 4) + 3n_1 n_2 \alpha + 3n_1 n_2 \alpha^2 + n_1(3n_1 + n - 4)\alpha^3] \\
& - v_3^{n-1} \frac{3b_{n-1} n_1 n_2}{8n(n-1)(n-2)} [n + 2n_2 - 4 + (n - 6n_2 + 4)\alpha \\
& + (n - 6n_1 + 4)\alpha^2 + (n + 2n_1 - 4)\alpha^3].
\end{aligned} \tag{51}$$

These four approximations for B_{n_1, n_2}^* satisfy the requirements (17) and (18). The condition (22) is only satisfied by the W approximation, while the condition (23) is only satisfied by the H approximation. By construction, the BS approximation, Eq. (51), reproduces the exact third virial coefficients (44). It is interesting to point out the approximations (46), (47) and (49) also yield the exact value of the third virial coefficients.

In order to assess the accuracy of the previous approximate expressions for B_{n_1, n_2}^* with $n \geq 4$, we will take the case $n_1 = 3$, $n_2 = 1$ as a benchmark for comparison since the coefficient $B_{3,1}^*$ has been obtained *exactly* for the interval $0 \leq \alpha \leq 2/\sqrt{3} - 1 \simeq 0.1547$ [17]. This exact result is

$$\begin{aligned}
B_{3,1}^* = & \frac{v_3^3}{4} \left(1 + 9\alpha + 36\alpha^2 + 21\alpha^3 + \frac{27}{2}\alpha^4 + \frac{27}{10}\alpha^5 \right. \\
& \left. - \frac{108}{5}\alpha^6 - \frac{648}{35}\alpha^7 - \frac{81}{14}\alpha^8 - \frac{9}{14}\alpha^9 \right).
\end{aligned} \tag{52}$$

Setting $n_1 = 3$ and $n_2 = 1$ in Eqs. (46), (47), (49) and (51), respectively, yields

$$B_{3,1}^{*SYH} = \frac{v_3^3}{4} \left[1 + \left(\frac{b_4}{2} + 1 \right) \alpha + (2b_4 - 5) \alpha^2 + 3 \left(\frac{b_4}{2} + 1 \right) \alpha^3 \right], \tag{53}$$

$$B_{3,1}^{*W} = \frac{v_3^3}{4} [1 + (28 - b_4) \alpha + (5b_4 - 59) \alpha^2 + 30\alpha^3], \tag{54}$$

$$B_{3,1}^{*H} = \frac{v_3^3}{4} [b_4 - 18 + 9\alpha + 36\alpha^2 + 3(b_4 - 9)\alpha^3] \tag{55}$$

and

$$B_{3,1}^{*BS} = \frac{v_3^3}{16} [b_4 - 15 + (3b_4 - 15)\alpha + (3b_4 + 75)\alpha^2 + (9b_4 - 45)\alpha^3]. \tag{56}$$

Note that Eqs. (53) and (54) would coincide if b_4 were equal to 18, as happens with the CS equation of state. However, they differ if we take the exact value $b_4 = 18.36477$. The maximum deviation of (54) from the exact result (52) is about 1.8% (at $\alpha \simeq 0.08$), while that of (53) is about 3.7% (at $\alpha \simeq 0.09$). Also, if b_4 were equal to 19, Eq. (55) would coincide with Eq. (54), while using the exact value for b_4 Eq. (55) implies that $\lim_{\alpha \rightarrow 0} B_{3,1}^{*H} = 0.091v_3^3$ instead of the exact value $\lim_{\alpha \rightarrow 0} B_{3,1}^* = \frac{1}{4}v_3^3$, which, however, is satisfied by the approximations (53) and (54). Comparison between Hamad's approximation (55) and the exact result (52) for $\alpha \leq 2/\sqrt{3} - 1$ shows that the relative error of (55) monotonically decreases from about 64% at $\alpha = 0$ to about 18% at $\alpha = 2/\sqrt{3} - 1$. Finally, according to Eq. (56) with the exact value of b_4 , $\lim_{\alpha \rightarrow 0} B_{3,1}^{*BS} = 0.210v_3^3$, while in the interval $\alpha \leq 2/\sqrt{3} - 1$ the relative error of (56) monotonically decreases from about 16% at $\alpha = 0$ to about 1.7% at $\alpha = 2/\sqrt{3} - 1$. This relative error is always larger than that of Eqs. (53) and (54), but smaller than that of Eq. (55).

Apart from the exact result for $B_{3,1}^*$, as pointed out before, numerical values of the coefficients B_{n_1, n_2}^* corresponding to $n \leq 5$ are also available for hard spheres [12–14,16]. In Figs. 11 and 12 we display the comparison between the theoretical estimates we have just discussed for the virial coefficients and the numerical results. It is clear that the estimates B_{n_1, n_2}^{*SYH} , B_{n_1, n_2}^{*W} and B_{n_1, n_2}^{*BS} for $n = 4$ and $n = 5$ are remarkably close to one another and to the exact results, except perhaps in the case of $B_{1,4}^{*BS}$. On the other hand, the prediction of B_{n_1, n_2}^{*H} is much poorer.




FIG. 11. Reduced composition-independent virial coefficients $B_{n_1, n-n_1}^*$ as functions of the size ratio α for $d = 3$, $n = 4$, and, from top to bottom, $n_1 = 1, 2$, and 3 . The circles are “exact” Monte Carlo results [13], the solid line is the theoretical prediction (46), the dashed line is the theoretical prediction (47), the dash-dot line is the theoretical prediction (49) and the dotted line is the theoretical prediction (51).




FIG. 12. Reduced composition-independent virial coefficients $B_{n_1, n-n_1}^*$ as functions of the size ratio α for $d = 3$, $n = 5$, and, from top to bottom, $n_1 = 1, 2, 3$ and 4 . The circles are “exact” Monte Carlo results [16,14], the solid line is the theoretical prediction (46), the dashed line is the theoretical prediction (47), the dash-dot line is the theoretical prediction (49) and the dotted line is the theoretical prediction (51).

Once we have briefly discussed the performance of the various approximate expressions for the composition-independent virial coefficients, we may next wonder how well the approximate EOS for a binary mixture of hard spheres perform with respect to the simulation data. To address this issue, we have first to specify what $Z_s(\eta)$ is. Two choices will be made here. On the one hand, since perhaps the most popular and widely known EOS for a single component fluid of hard spheres is the CS EOS, we will also use it here. On the other hand, a more accurate EOS for the same system is obtained if a rescaled Padé approximation taking into account the known virial coefficients is used [24,44]. In Tables III and IV we present a comparison of the results obtained using the various approximate EOS for the mixture with these two choices of $Z_s(\eta)$, two different mole fractions and a diameter ratio $\alpha = 0.3$. In Table III, the values corresponding to the widely used Boublík-Mansoori-Carnahan-Starling-Leland (BMCSL) EOS [45] have also been included. One immediately notes that, as already stated in the literature [24,32,35], the EOS given in Eq. (2) using the CS EOS for the single component system provides a very accurate account of the simulation data, better than any of the other EOS for mixtures considered, including that of Eq. (39). If instead of the CS EOS the rescaled Padé approximation is taken for $Z_s(\eta)$, then it is Eq. (39) which gives in general the best results, although the original recipe (2) and the BS EOS (43) are not too far behind, the latter being especially accurate in the case $x_1 = 0.75$. It should be noted that, irrespective of the choice for $Z_s(\eta)$, both $Z_m^H(\eta)$ and $Z_m^{BS}(\eta)$ for these size ratio and packing fractions always underestimate the simulation results, a comment that also applies to the BMCSL EOS.

TABLE III. Comparison of the compressibility factor from several equations of state with simulation values (Z_{simul}) for binary additive mixtures of hard spheres ($d = 3$) with a size ratio $\alpha = 0.3$. The fourth to seventh columns give the deviations from Z_{simul} of the values of the compressibility factor as obtained from the equations of state (2), (39), (42) and (43), respectively. In these equations of state, the Carnahan-Starling compressibility factor for the one-component fluid [25] has been used. For the sake of comparison, also the results arising from the Boublík-Mansoori-Carnahan-Starling-Leland (BMCSL) equation of state [45] are included.

x_1	η	$Z_{\text{simul}}^{\text{a}}$	Eq. (2) ^b	Eq. (39) ^c	Eq. (42) ^d	Eq. (43) ^e	BMCSL
0.0625	0.3	2.790	-0.001	-0.009	-0.061	-0.016	-0.014
	0.35	3.473	0.006	-0.011	-0.112	-0.024	-0.020
	0.4	4.410	0.013	-0.019	-0.210	-0.043	-0.035
	0.45	5.722	0.027	-0.033	-0.385	-0.076	-0.063
	0.49	7.158	0.065	-0.036	-0.603	-0.102	-0.081
0.75	0.3	3.554	-0.005	-0.006	-0.024	-0.010	-0.008
	0.35	4.601	-0.012	-0.014	-0.048	-0.021	-0.018
	0.4	6.045	-0.010	-0.014	-0.079	-0.027	-0.021
	0.45	8.097	-0.002	-0.011	-0.129	-0.033	-0.022
	0.49	10.415	-0.004	-0.021	-0.210	-0.056	-0.037

^aRef. [46]

^bRef. [32]

^cThis work

^dRef. [40]

^eRef. [43]

TABLE IV. Comparison of the compressibility factor from several equations of state with simulation values (Z_{simul}) for binary additive mixtures of hard spheres ($d = 3$) with a size ratio $\alpha = 0.3$. The fourth to seventh columns give the deviations from Z_{simul} of the values of the compressibility factor as obtained from the equations of state (2), (39), (42) and (43), respectively. In these equations of state, a rescaled Padé approximant compressibility factor for the one-component fluid [24,44] has been used.

x_1	η	$Z_{\text{simul}}^{\text{a}}$	Eq. (2) ^b	Eq. (39) ^c	Eq. (42) ^d	Eq. (43) ^e
0.0625	0.3	2.790	0.004	-0.005	-0.050	-0.010
	0.35	3.473	0.014	-0.004	-0.095	-0.014
	0.4	4.410	0.025	-0.008	-0.186	-0.029
	0.45	5.722	0.043	-0.018	-0.353	-0.058
	0.49	7.158	0.084	-0.015	-0.564	-0.081
0.75	0.3	3.554	0.003	0.002	-0.013	-0.001
	0.35	4.601	0.001	-0.001	-0.032	-0.007
	0.4	6.045	0.010	0.006	-0.055	-0.007
	0.45	8.097	0.025	0.016	-0.096	-0.006
	0.49	10.415	0.028	0.012	-0.171	-0.023

^aRef. [46]

^bRef. [32]

^cThis work

^dRef. [40]

^eRef. [43]

V. CONCLUDING REMARKS

In this paper we have further assessed the merits and limitations of a simple recipe recently introduced [32] to derive the compressibility factor of a multicomponent mixture of d -dimensional hard spheres. We have now considered the case of binary mixtures and looked in particular at the composition-independent virial coefficients. The comparison with the available exact and simulation results confirmed the usefulness of our approach except perhaps when the mixture involves components of very disparate sizes (especially for high dimensionalities). Guided by the conditions that the virial coefficients must fulfill in certain limits, a slight modification of the form of the original coefficients was made, trying to sacrifice simplicity as less as possible while at the same time improving the accuracy of their numerical predictions. It is fair to state that the (modified) composition-independent virial coefficients are in excellent agreement with the reported values for all dimensions and all size ratios.

Due to its particular relevance, the case of the binary mixture of hard spheres ($d = 3$) was analyzed in some detail, both in connection with the virial coefficients and through the consideration of different EOS for this system that exist in the literature. As for the virial coefficients, the trends pointed out above about the reliability of the original and modified prescriptions hold also for $d = 3$. The comparison with the simulation results for the compressibility factor of these mixtures given in Tables III and IV indicates the superiority of the original recipe for the EOS provided one uses the CS EOS for $Z_s(\eta)$. On the other hand, if the more accurate rescaled Padé approximation for $Z_s(\eta)$ is used, it is the new EOS (39) (i.e., the one obtained after resumming the virial series with the modified coefficients) which gives the best performance, although only slightly better than $Z_m(\eta)$ as given by Eq. (2) with $d = 3$. Both equations are also of comparable accuracy to the one of the presumably best EOS available presently for hard-sphere mixtures, namely the rescaled Padé approximant for mixtures introduced by Malijevský and Veverka [24]. This latter equation of state rests on a different philosophy, however, in the sense that it makes use of the known virial coefficients of the mixture rather than the $Z_s(\eta)$, as in our approach and those of Hamad [40] and Barrio and Solana [43].

In conclusion, we want to add that both the original proposal of the EOS of a binary mixture of d -dimensional hard spheres, Eq. (2), and the modified version, Eq. (34), provide two simple and yet accurate EOS at least with respect to the composition-independent virial coefficients and in comparison with the (relatively scarce) simulation data. Further confirmation of their usefulness depends on new simulations which we hope will be encouraged by the results presented in this paper.

ACKNOWLEDGMENTS

A.S. and S.B.Y. acknowledge partial support from the DGES (Spain) through grant No. PB97-1501 and from the Junta de Extremadura (Fondo Social Europeo) through grant No. IPR99C031. They are also grateful to the DGES (Spain) for sabbatical grants No. PR2000-0117 and No. PR2000-0116, respectively. The research of M.L.H. was supported in part by DGAPA-UNAM under Project IN103100.

APPENDIX A: DERIVATION OF EQS. (7)–(10)

In order to identify the composition-independent coefficients B_{n_1, n_2} from Eq. (5), we need to expand the moments $\langle \sigma^p \rangle$. After some algebra, we get

$$\begin{aligned} \langle \sigma^d \rangle^{n-3} \langle \sigma^{d-m} \rangle \langle \sigma^{m+1} \rangle \langle \sigma^{d-1} \rangle &= \sigma_1^{d(n-1)} \sum_{n_1=0}^n \alpha^{d(n-n_1-1)} \frac{(n-3)!}{n_1!(n-n_1)!} x_1^{n_1} x_2^{n-n_1} \\ &\times [n_1(n_1-1)(n_1-2)\alpha^d + n_1(n_1-1)(n-n_1)\alpha^{d-1} \\ &+ n_1(n_1-1)(n-n_1)\alpha^{d-m} + n_1(n-n_1)(n-n_1-1)\alpha^{d-m-1} \\ &+ n_1(n_1-1)(n-n_1)\alpha^{m+1} + n_1(n-n_1)(n-n_1-1)\alpha^m \\ &+ n_1(n-n_1)(n-n_1-1)\alpha \\ &+ (n-n_1)(n-n_1-1)(n-n_1-2)], \end{aligned} \quad (\text{A1})$$

$$\begin{aligned} \langle \sigma^d \rangle^{n-2} \langle \sigma^{d-m} \rangle \langle \sigma^m \rangle &= \sigma_1^{d(n-1)} \sum_{n_1=0}^n \alpha^{d(n-n_1-1)} \frac{(n-2)!}{n_1!(n-n_1)!} x_1^{n_1} x_2^{n-n_1} \\ &\times [n_1(n_1-1)\alpha^d + n_1(n-n_1)\alpha^{d-m}] \end{aligned}$$

$$+n_1(n-n_1)\alpha^m + (n-n_1)(n-n_1-1)], \quad (\text{A2})$$

where $\alpha \equiv \sigma_2/\sigma_1$. Now we use the identities

$$\sum_{m=0}^d \frac{d!}{m!(d-m)!} \alpha^m = (1+\alpha)^d, \quad (\text{A3})$$

$$\sum_{m=0}^d \frac{(d-1)!}{m!(d-m)!} (d-m)\alpha^m = (1+\alpha)^{d-1}, \quad (\text{A4})$$

to get

$$\begin{aligned} & \sum_{m=0}^d \frac{(d-1)!}{m!(d-m)!} (d-m) \langle \sigma^d \rangle^{n-3} \langle \sigma^{d-m} \rangle \langle \sigma^{m+1} \rangle \langle \sigma^{d-1} \rangle = \sigma_1^{d(n-1)} \sum_{n_1=0}^n \alpha^{d(n-n_1-1)} \frac{(n-3)!}{n_1!(n-n_1)!} \\ & \times x_1^{n_1} x_2^{n-n_1} [n_1(n_1-1)(n_1-2)2^{d-1}\alpha^d + n_1(n_1-1)(n-n_1)2^{d-1}\alpha^{d-1} \\ & + 2n_1(n_1-1)(n-n_1)\alpha(1+\alpha)^{d-1} + 2n_1(n-n_1)(n-n_1-1)(1+\alpha)^{d-1} \\ & + n_1(n-n_1)(n-n_1-1)2^{d-1}\alpha + (n-n_1)(n-n_1-1)(n-n_1-2)2^{d-1}], \end{aligned} \quad (\text{A5})$$

$$\begin{aligned} & \sum_{m=0}^d \frac{d!}{m!(d-m)!} \langle \sigma^d \rangle^{n-2} \langle \sigma^{d-m} \rangle \langle \sigma^m \rangle = \sigma_1^{d(n-1)} \sum_{n_1=0}^n \alpha^{d(n-n_1-1)} \frac{(n-2)!}{n_1!(n-n_1)!} x_1^{n_1} x_2^{n-n_1} \\ & \times [n_1(n_1-1)2^d\alpha^d + 2n_1(n-n_1)(1+\alpha)^d + (n-n_1)(n-n_1-1)2^d]. \end{aligned} \quad (\text{A6})$$

From here we finally obtain Eqs. (7)–(10).

APPENDIX B: DERIVATION OF EQS. (34)–(38)

Let us rewrite Eq. (7) as

$$\begin{aligned} B_{n_1, n_2} = v_d^{n-1} & \left[C_{n_1, n_2}^{(1)} \sigma_1^{d(n_1-1)} \sigma_2^{dn_2} + C_{n_1, n_2}^{(2)} \sigma_1^{d(n_1-1)+1} \sigma_2^{dn_2-1} + C_{n_1, n_2}^{(3)} 2^{d-1} \sigma_{12}^{d-1} \sigma_1^{d(n_1-1)+1} \sigma_2^{d(n_2-1)} \right. \\ & \left. + 1 \leftrightarrow 2 \right], \end{aligned} \quad (\text{B1})$$

where $\sigma_{12} \equiv (\sigma_1 + \sigma_2)/2$ and the notation $1 \leftrightarrow 2$ indicates that a contribution similar to the three summands appearing in the brackets in which the roles of (n_1, n_2) and of (σ_1, σ_2) are interchanged (i.e., $n_1 \rightarrow n_2$, $\sigma_1 \rightarrow \sigma_2$ and vice versa) should be added. The compressibility factor of the mixture is then given by

$$\begin{aligned} Z_m(\eta) - 1 &= \sum_{n=2}^{\infty} \rho^{n-1} \sum_{n_1=0}^n B_{n_1, n_2} \frac{n!}{n_1!n_2!} x_1^{n_1} x_2^{n_2} \\ &= \sum_{n=2}^{\infty} (v_d \rho)^{n-1} \sum_{n_1=0}^n \frac{n!}{n_1!n_2!} x_1^{n_1} x_2^{n_2} \left[C_{n_1, n_2}^{(1)} \sigma_1^{d(n_1-1)} \sigma_2^{dn_2} + C_{n_1, n_2}^{(2)} \sigma_1^{d(n_1-1)+1} \sigma_2^{dn_2-1} \right. \\ & \quad \left. + C_{n_1, n_2}^{(3)} 2^{d-1} \sigma_{12}^{d-1} \sigma_1^{d(n_1-1)+1} \sigma_2^{d(n_2-1)} \right] \\ & \quad + 1 \leftrightarrow 2. \end{aligned} \quad (\text{B2})$$

Now we insert Eqs. (27)–(30) and sum over n_1 and n . To do that, we use the identities

$$x_1 [Z_s(\eta) - 1] = \sum_{n=2}^{\infty} (v_d \rho)^{n-1} \sum_{n_1=0}^n \frac{n!}{n_1!n_2!} x_1^{n_1} x_2^{n_2} b_n \frac{n_1}{n} \sigma_1^{d(n_1-1)} \sigma_2^{dn_2}, \quad (\text{B3})$$

$$x_2 [Z_s(\eta) - 1] = \sum_{n=2}^{\infty} (v_d \rho)^{n-1} \sum_{n_1=0}^n \frac{n!}{n_1!n_2!} x_1^{n_1} x_2^{n_2} b_n \frac{n_2}{n} \sigma_1^{dn_1} \sigma_2^{d(n_2-1)}, \quad (\text{B4})$$

$$\langle \sigma^d \rangle [Z_s(\eta) - 1] = \sum_{n=2}^{\infty} (v_d \rho)^{n-1} \sum_{n_1=0}^n \frac{n!}{n_1! n_2!} x_1^{n_1} x_2^{n_2} b_n \sigma_1^{dn_1} \sigma_2^{dn_2}, \quad (\text{B5})$$

$$x_1 x_2 \frac{Z_s(\eta) - 1}{\langle \sigma^d \rangle} = \sum_{n=2}^{\infty} (v_d \rho)^{n-1} \sum_{n_1=0}^n \frac{n!}{n_1! n_2!} x_1^{n_1} x_2^{n_2} b_n \frac{n_1 n_2}{n(n-1)} \sigma_1^{d(n_1-1)} \sigma_2^{d(n_2-1)}, \quad (\text{B6})$$

$$x_1 x_2 \frac{\eta}{(1-\eta) \langle \sigma^d \rangle} = \sum_{n=2}^{\infty} (v_d \rho)^{n-1} \sum_{n_1=0}^n \frac{n!}{n_1! n_2!} x_1^{n_1} x_2^{n_2} \frac{n_1 n_2}{n(n-1)} \sigma_1^{d(n_1-1)} \sigma_2^{d(n_2-1)}. \quad (\text{B7})$$

In addition, [37]

$$\frac{x_1}{1-\eta_2} Z_s \left(\frac{\eta_1}{1-\eta_2} \right) - x_1 = \sum_{n=2}^{\infty} (v_d \rho)^{n-1} \sum_{n_1=0}^n \frac{n!}{n_1! n_2!} x_1^{n_1} x_2^{n_2} b_{n_1} \frac{n_1}{n} \sigma_1^{d(n_1-1)} \sigma_2^{dn_2}, \quad (\text{B8})$$

$$\frac{x_2}{1-\eta_1} Z_s \left(\frac{\eta_2}{1-\eta_1} \right) - x_2 = \sum_{n=2}^{\infty} (v_d \rho)^{n-1} \sum_{n_1=0}^n \frac{n!}{n_1! n_2!} x_1^{n_1} x_2^{n_2} b_{n_2} \frac{n_2}{n} \sigma_1^{dn_1} \sigma_2^{d(n_2-1)}. \quad (\text{B9})$$

This allows one to obtain $Z_m(\eta)$ in terms of Z_s evaluated at η , $\eta_1/(1-\eta_2)$ and $\eta_2/(1-\eta_1)$. After some algebra, we arrive at Eqs. (34)–(38).

* Permanent address: Departamento de Física, Universidad de Extremadura, E-06071 Badajoz, Spain; Email address: andres@unex.es

† Permanent address: Departamento de Física, Universidad de Extremadura, E-06071 Badajoz, Spain; Email address: santos@unex.es

‡ Also Consultant at Programa de Simulación Molecular del Instituto Mexicano del Petróleo; Email address: malopez@servidor.unam.mx

- [1] VAN DER WAALS, J. D., *Ph. D. dissertation*, 1873, Leiden. Translated in ROWLINSON, J. S., 1988, *Studies in Statistical Mechanics*, vol. XIV (Amsterdam: North-Holland).
- [2] KAMMERLINGH ONNES, H., 1909, *Comm. Leiden*, **71**.
- [3] See, for instance, MASON, E. A., AND SPURLING, T. H., 1969, *The virial equation of state* (New York: Pergamon).
- [4] MAYER, J. E., AND MAYER, M. G., 1940, *Statistical Mechanics* (New York: Wiley), Chapter 13.
- [5] Early analytical results on the virial coefficients of hard-sphere fluids ($d = 3$) up to the fourth include JÄGGER, G., 1896, *Sitzber. Akad. Wiss. Wien, Math. Natur.-w. Kl.* (Pt. 2a), **105**, 15, and BOLTZMANN, L., 1896, *Sitzber. Akad. Wiss. Wien, Math. Natur.-w. Kl.* (Pt. 2a), **105**, 695; 1899, *Proc. Kon. Acad. V. Wetensch., Amsterdam*, **1**, 398. Those of hard-disc fluids ($d = 2$) are due to TONKS, L., 1936, *Phys. Rev.*, **50**, 955; ROWLINSON, J. S., 1963, *Molec. Phys.*, **7**, 593, and HEMMER, P. C., 1965, *J. chem. Phys.*, **42**, 1116. Numerical calculations of higher coefficients up to $n = 7$ for both hard discs and hard spheres can be found in REE, F. H., AND HOOVER, W. G., 1967, *J. chem. Phys.*, **46**, 4181; KRATKY, K. W., 1976, *Physica A*, **85**, 607; 1977, *Physica A*, **87**, 584; 1982, *J. statist. Phys.*, **27**, 533; 1982, *J. statist. Phys.*, **29**, 129, and JANSE VAN RENSBURG, E. J., AND TORRIE, G. M., 1992, *J. Phys. A*, **26**, 943. The most recent calculations for these dimensions up to $n = 8$ are those of Ref. [6]. In the case of hard hyperspheres in 4D and 5D, the available results are those of Refs. [7] and [8].
- [6] JANSE VAN RENSBURG, E. J., 1993, *J. Phys. A*, **26**, 4805.
- [7] LUBAN, M., AND BARAM, A., 1982, *J. chem. Phys.*, **76**, 3233.
- [8] JOSLIN, C. J., 1982, *J. chem. Phys.*, **77**, 2701.
- [9] KIHARA, T., AND MIYOSHI, K., 1975, *J. statist. Phys.*, **13**, 337.
- [10] BARRAT, J.-L., XU, H., HANSEN, J.-P., AND BAUS, M., 1988, *J. Phys. C*, **21**, 3165.
- [11] SAIJA, F., FIUMARA, G., AND GIAQUINTA, P. V., 1996, *Molec. Phys.*, **87**, 991.
- [12] ENCISO, E., ALMARZA, N. G., CALZAS, D. S., AND GONZÁLEZ, M. A., 1997, *Molec. Phys.*, **92**, 173.
- [13] SAIJA, F., FIUMARA, G., AND GIAQUINTA, P. V., 1997, *Molec. Phys.*, **92**, 1089.

- [14] ENCISO, E., ALMARZA, N. G., GONZÁLEZ, M. A., AND BERMEJO, F. J., 1998, *Phys. Rev. E*, **57**, 4486.
- [15] WHEATLEY, R. J., 1998, *Molec. Phys.*, **93**, 675.
- [16] WHEATLEY, R. J., SAIJA, F., AND GIAQUINTA, P. V., 1998, *Molec. Phys.*, **94**, 877.
- [17] BLAAK, R., 1998, *Molec. Phys.*, **95**, 695.
- [18] WHEATLEY, R. J., 1999, *Molec. Phys.*, **96**, 1805.
- [19] ENCISO, E., ALMARZA, N. G., GONZÁLEZ, M. A., AND BERMEJO, F. J., 2001, unpublished. We want to thank these authors for providing their results to us prior to publication.
- [20] ERPENBECK, J. J., AND WOOD, W. W., 1984, *J. statist. Phys.*, **35**, 321.
- [21] ERPENBECK, J. J., AND LUBAN, M., 1985, *Phys. Rev. A*, **32**, 2920.
- [22] SANCHEZ, I. C., 1994, *J. chem. Phys.*, **101**, 7003.
- [23] COUSSAERT, T., AND BAUS, M., 1998, *J. chem. Phys.*, **109**, 6012.
- [24] MALIJEVSKÝ, A., AND VEVERKA, J., 1999, *Phys. Chem. Chem. Phys.*, **1**, 4267.
- [25] CARNAHAN, N. F., AND STARLING, K. E., 1969, *J. chem. Phys.*, **51**, 635.
- [26] Particular examples for hard discs are discussed in SANTOS, A., LÓPEZ DE HARO, M., AND YUSTE, S. B., 1995, *J. chem. Phys.*, **103**, 4622. The most important equations of state for hard spheres have been recently examined in Ref. [24].
- [27] BAUS, M., AND COLOT, J. L., 1987, *Phys. Rev. A*, **36**, 3912.
- [28] SONG, Y., AND MASON, E. A., 1990, *J. chem. Phys.*, **93**, 686.
- [29] LUBAN, M., AND MICHELS, J. P. J., 1990, *Phys. Rev. A*, **41**, 6796.
- [30] BISHOP, M., MASTERS, A., AND CLARKE, J. H. R., 1999, *J. chem. Phys.*, **110**, 11449.
- [31] SANTOS, A., 2000, *J. chem. Phys.*, **112**, 10680.
- [32] SANTOS, A., YUSTE, S. B., AND LÓPEZ DE HARO, M., 1999, *Molec. Phys.*, **96**, 1.
- [33] WHEATLEY, R. J., 1998, *Molec. Phys.*, **93**, 965.
- [34] WHEATLEY, R. J., 1999, *J. chem. Phys.*, **111**, 5455.
- [35] CAO, D., CHAN, K.-Y., HENDERSON, D. AND WANG, W., 2000, *Molec. Phys.*, **98**, 619.
- [36] GONZÁLEZ MELCHOR, M., ALEJANDRE, J., AND LÓPEZ DE HARO, M., 2001, *J. chem. Phys.*, **114**, 4905.
- [37] SANTOS, A., 1999, *Molec. Phys.*, **96**, 1185. [Erratum: 2001, *Molec. Phys.*, **99**, 617.]
- [38] TUKUR, N. M., HAMAD, E. Z., AND MANSOORI, G. A., 1999, *J. chem. Phys.*, **110**, 3463.
- [39] REISS, H., FRISCH, H. L., AND LEBOWITZ, J. L., 1959, *J. chem. Phys.*, **31**, 369; HELFAND, E., FRISCH, H. L., AND LEBOWITZ, J. L., 1961, *J. chem. Phys.*, **34**, 1037.
- [40] HAMAD, E. Z., 1995, *J. chem. Phys.*, **103**, 3733.
- [41] HAMAD, E., 1994, *J. chem. Phys.*, **101**, 10195.
- [42] For additional consistency conditions on the contact values of g_{ij} , see BARRIO, C., AND SOLANA, J. R., 2000, *J. chem. Phys.*, **113**, 10180.
- [43] BARRIO, C., AND SOLANA, J. R., 1999, *Molec. Phys.*, **97**, 797. For the adaptation of this procedure to the case of hard-disk mixtures, see BARRIO, C., AND SOLANA, J. R., 2001, *Phys. Rev. E*, **63**, 011201.
- [44] We want to point out that the approximation used in Ref. [24] is somewhat different to the one used here, since apparently the number of significative figures in the value of the virial coefficients is not the same in both instances. We have checked that the present approximation yields practically identical results.
- [45] BOUBLÍK, T., 1970, *J. chem. Phys.*, **53**, 471; MANSOORI, G. A., CARNAHAN, N. F., STARLING, K. E., AND LELAND, T. W., 1971, *J. chem. Phys.*, **54**, 1523.
- [46] BAROŠOVÁ, A., MALIJEVSKÝ, A., LABÍK, S., AND SMITH, W. R., 1996, *Molec. Phys.*, **87**, 423.